



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$x = \frac{a^2 - b^2}{y} = \frac{500}{y} = 35.559825. \quad x + y = 49.620636, \quad x - y = 21.499014.$$

$$AB = \sqrt{(AC^2 - BC^2)} = \sqrt{[a^2 - (x+y)^2]} = 33.731178.$$

Solutions were also received from G. B. M. ZERR, J. F. W. SCHEFFER, L. B. FRAKER, and J. K. ELLWOOD.

25. Proposed by L. B. FRAKER, Weston, Ohio.

The sides of a quadrilateral board are $AB=7$ inches, $BC=15$ inches, $CD=21$ inches, and $DA=13$ inches; radius of incircled circle is 6 inches. (1) What are dimensions of the largest rectangular board that can be cut out of the given board, (2) largest square, (3) largest equilateral triangle? (Please solve without use of the calculus.)

Solution by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

Let $Aa=Ad=x$, $Ba=Bb=y$, $Cb=Cc=z$, $Dc=Dd=w$,

$$\text{then } x+y=AB=7=6 \cot \frac{1}{2}A + 6 \cot \frac{1}{2}B \dots (1),$$

$$y+z=BC=15=6 \cot \frac{1}{2}B + 6 \cot \frac{1}{2}C \dots (2),$$

$$z+w=CD=21=6 \cot \frac{1}{2}C + 6 \cot \frac{1}{2}D \dots (3),$$

$$w+x=DA=13=6 \cot \frac{1}{2}D + 6 \cot \frac{1}{2}A \dots (4).$$

$$\therefore 28=6 \cot \frac{1}{2}A + 6 \cot \frac{1}{2}B + 6 \cot \frac{1}{2}C + 6 \cot \frac{1}{2}D \dots (5).$$

$$(5) - \{ (1) + (4) \} \text{ gives } 4 = 3 \cot \frac{1}{2}C - 3 \cot \frac{1}{2}A$$

$$\therefore 4 \sin \frac{1}{2}A \sin \frac{1}{2}C = 3 \sin \frac{1}{2}A \cos \frac{1}{2}C - 3 \cos \frac{1}{2}A \sin \frac{1}{2}C \dots (6).$$

$$(5) - \{ (3) + (4) \} \text{ gives } 1 = \cos \frac{1}{2}D - \cos \frac{1}{2}B$$

$$\therefore \sin \frac{1}{2}D \sin \frac{1}{2}B = \sin \frac{1}{2}B \cos \frac{1}{2}D - \cos \frac{1}{2}B \sin \frac{1}{2}D \dots (7).$$

From DB^2 we get $45 \cos C - 13 \cos A = 32$.

$$\therefore 13 \sin^2 \frac{1}{2}A = 45 \sin^2 \frac{1}{2}C \dots (8).$$

$$45 \cos^2 \frac{1}{2}C - 13 \cos^2 \frac{1}{2}A = 32 \dots (9).$$

From AC^2 we get $13 \cos D - 5 \cos B = 8$.

$$\therefore 13 \sin^2 \frac{1}{2}D = 5 \sin^2 \frac{1}{2}B \dots (10).$$

$$13 \cos^2 \frac{1}{2}D - 5 \cos^2 \frac{1}{2}B = 8 \dots (11).$$


From (6), (7), (8), (9), (10), (11) we get

$$A=134^\circ 45' 35'', \quad B=106^\circ 15' 39'', \quad C=D=59^\circ 29' 23''.$$

Now the greatest equilateral triangle is the one which has the greatest side. Since $D=C$ only two cases confront us, the one is to draw a line from B making an angle of 60° with DC , the other to draw a line from some point in DC making an angle of 60° with BC . The former line is found to be very little the longer. Let BL be this line, then we have $15 : BL = \sin 120^\circ : \sin 59^\circ 29' 23''$.

$\therefore BL = \text{side of triangle} = 14.92$ inches. $\therefore BLM = \text{triangle}$. Also since $D=C$, the greatest square has its side coincident with DC . Hence $\frac{21-x}{2} : x = \cos 59^\circ 29' 23'' : \sin 59^\circ 29' 23''$.

$\therefore x = \text{side of square} = 9.64$ inches, and $PQRS = \text{square}$. Also since $D=C$, the side of the greatest rectangle will



coincide with DC . Draw AI parallel and AK perpendicular to DC and let $EFGH$ be the rectangle.

Then $\frac{1}{2}(AI+DC) \times AK = EF \times FG + FG \times GC + \frac{1}{2}(AI+EF)(AK-KR')$.
But $AK=11.2$ inches, $AI=7.8$ inches. $\therefore 588=28 EF+33 FG$.

\therefore for maximum 28 $EF=33 FG$. $\therefore EF=10.5$ inches, FG 8.91 inches.

26. Proposed by J. F. W. SCHEFFER, A. M., Hagerstown, Maryland.

$ABCD$ represents a triangle, and $ABEF$ a trapezoid which is perpendicular to the rectangle, both figures having the side AB common to each other, and ADF and BCE forming two right triangles perpendicular to the rectangle $ABCD$. To determine the conoidal surface $CDFE$ so as to satisfy the condition that any plane laid through AB will intersect it in a straight line. Also find volume of the surface thus formed.

Solution by the PROPOSER.

Let $BC=AD=h$, $AB=a$, $BE=b$, $AF=c$. Let P represent a point in the surface, and put $AR=x$, $RQ=y$, $PQ=z$

The triangles BGK , PQR , and AHI are similar, and we may now put $AH=ny$, $IH=nz$, $BG=my$, $KG=mz$; but $h:mz=b-my$; $h:nz=c:c-nz$, whence $m=\frac{bh}{hy+bz}$, $n=\frac{ch}{hy+cz}$.

In the trapezoid $AHGB$, we now have
 $AB=a$, $AH=\frac{chy}{hy+cz}$, $BG=\frac{bhy}{hy+bz}$, $AR=x$,

$RQ=y$. $\therefore (AH+y)x + (BG+y)(a-x) = (AH+BG)a$.

Substituting, clearing of fractions, and arranging, we find for the equation of the surface

$$abcz^2 + a(b+c)hyz - (b-c)h^2xy + ah^2y^2 - abchz - ach^2y = 0.$$

Let us now denote $\angle CBK = \angle DAI$ by θ , and angles BCK and ADI represent by C and D . For the volume we have $\frac{1}{6}ah^2 \int_0^{\pi} \left[\frac{\sin^2 C}{\sin^2(C+\theta)} \right.$

$$+ \frac{\sin^2 D}{\sin^2(D+\theta)} + \frac{\sin C \sin D}{\sin(C+\theta) \sin(D+\theta)} \Big] d\theta = \frac{1}{6}ah^2 \left[\tan C + \tan D \right. \\ \left. + \frac{\tan C \tan D}{\tan C - \tan D} \log \frac{\tan C}{\tan D} \right] \text{ but } \tan C = \frac{b}{h}, \tan D = \frac{c}{h};$$

$$\therefore \text{volume} = \frac{1}{6}ah \left[b+c + \frac{bc}{b-c} \log \frac{b}{c} \right].$$

27. Proposed by ADOLPH BAILOFF, Durand Wisconsin.

A line BF , that bisects an angle exterior to the vertical angle of an isosceles triangle is parallel to the base AC .

Solution by Mrs. MARY E. HOGSETT, Danville, Kentucky; P. S. BERG, Apple Creek, Ohio, Professors JOHN FAUGHT, Bloomington, Indiana; and M. A. GRUBER, War Department, Washington, D. C.

